

Cognitive limitations and diabetes comorbidity among racial and ethnically diverse older adults by nativity status in the United States

Tiffany B. Kindratt,¹ Berford L. Moncriffe¹

¹University of Texas at Arlington



INTRODUCTION

- Diabetes is one of the most common comorbid conditions among adults with Alzheimer's disease and related dementias in the U.S.
- Disparities in the prevalence of cognitive limitations and diabetes by race, ethnicity, and nativity status have been identified
- Few studies have examined the prevalence of having both cognitive limitations and diabetes among US and foreign-born racial and ethnic groups

OBJECTIVE

- 1) Estimate and compare the prevalence of cognitive limitations and diabetes among US and foreign-born White, Black, and Hispanic adults
- 2) Examine differences after adjusting for modifiable risk factors

METHODS

Sample and Participants

- Linked 2000-2017 National Health Interview Survey (NHIS) and 2001-2018 Medical Expenditure Panel Survey (MEPS)
- Total number of adults ages 65+ years (n=35,849)

Independent variable: Combined variable describing race, ethnicity and nativity status (3 groups)

- Foreign-born and US-born non-Hispanic Whites
- Foreign-born and US-born non-Hispanic Blacks
- Foreign-born and US born Hispanics

Dependent variable: Any cognitive limitation (yes/no) measured based on "yes" responses to:

- "experience confusion or memory loss"
- "have problems making decisions to the point that it interferes with daily activities"
- "require supervisions for their own safety"

Covariates:

- Age, sex, <9th grade level of education
- ADRD risk factors that are potentially modifiable in middle or late life (hearing loss, hypertension, obesity, smoking, physical inactivity, social isolation, diabetes)

Analysis

- Age and sex-adjusted prevalence estimates, and multivariable logistic regression models calculated using STATA

RESULTS

Table 1: Crude and adjusted logistic regression results, 2000-2017 NHIS and 2001-2018 MEPS, n=35,849

	Model 1 ^a	Model 2 ^b	Model 3 ^c	Model 4 ^d
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
US-born NH white (reference)	1.00	1.00	1.00	1.00
US-born NH Black	2.30 (1.94, 2.72)	2.43 (2.06, 2.87)	2.22 (1.86, 2.66)	1.94 (1.58, 2.39)
US-born Hispanic	2.24 (1.72, 2.92)	2.41 (1.84, 3.16)	1.87 (1.39, 2.51)	1.97 (1.41, 2.76)
Foreign-born NH white	1.56 (1.00, 2.42)	1.46 (0.94, 2.26)	1.36 (0.87, 2.14)	1.36 (0.80, 2.32)
Foreign-born NH black	2.24 (1.51, 3.32)	2.48 (1.66, 3.70)	2.05 (1.35, 3.11)	2.44 (1.48, 4.02)
Foreign-born Hispanic	2.46 (1.97, 3.07)	2.61 (2.09, 3.26)	1.77 (1.36, 2.30)	1.64 (1.24, 2.18)
US-born NH Black (reference)	1.00	1.00	1.00	1.00
US-born NH White	0.43 (0.37, 0.51)	0.41 (0.35, 0.49)	0.45 (0.38, 0.54)	0.51 (0.42, 0.63)
US-born Hispanic	0.97 (0.73, 1.29)	0.99 (0.75, 1.32)	0.84 (0.62, 1.13)	1.01 (0.71, 1.44)
Foreign-born NH white	0.68 (0.43, 1.05)	0.60 (0.39, 0.93)	0.61 (0.39, 0.96)	0.70 (0.41, 1.20)
Foreign-born NH black	0.97 (0.65, 1.45)	1.02 (0.68, 1.53)	0.92 (0.61, 1.40)	1.26 (0.76, 2.07)
Foreign-born Hispanic	1.07 (0.84, 1.36)	1.07 (0.84, 1.36)	0.80 (0.62, 1.03)	0.85 (0.65, 1.10)
US-born Hispanic (reference)	1.00	1.00	1.00	1.00
US-born NH White	0.45 (0.34, 0.58)	0.41 (0.32, 0.54)	0.53 (0.40, 0.72)	0.51 (0.36, 0.71)
US-born NH Black	1.03 (0.77, 1.36)	1.01 (0.76, 1.34)	1.19 (0.88, 1.60)	0.98 (0.69, 1.40)
Foreign-born NH white	0.69 (0.42, 1.14)	0.61 (0.37, 0.99)	0.73 (0.44, 1.21)	0.69 (0.39, 1.23)
Foreign-born NH black	1.00 (0.63, 1.58)	1.03 (0.65, 1.63)	1.10 (0.68, 1.77)	1.24 (0.69, 2.21)
Foreign-born Hispanic	1.10 (0.78, 1.54)	1.08 (0.77, 1.51)	0.95 (0.67, 1.34)	0.83 (0.56, 1.24)


^aUnadjusted model

^bAdjusted for age and sex

^cAdjusted for Model 2+early life risk factor (highest level of education)

^dAdjusted for Model 3+selected risk factors measured during late life (hearing loss, hypertension, depression, social isolation, diabetes)

Figure 1: Prevalence of diabetes among adults with cognitive limitations (US born)



DISCUSSION

- For all groups, the odds ratio of having diabetes and cognitive limitations is highest among foreign-born NH Blacks (OR =2.44; 95% CI = 1.48, 4.02)
- The prevalence of diabetes and cognitive limitations among US born adults was highest among Hispanics (47.2%; SE=0.02)
- Results suggest a large potential burden of comorbid diabetes and cognitive limitations in US and foreign-born older adults
- Further studies will explore prevalence of other comorbidities and cognitive limitations

ACKNOWLEDGMENTS

- Funded by the Michigan Center for Contextual Factors in Alzheimer's Disease (MCCFAD) P30AG05930001, National Institute on Aging
- This research was conducted at the Dallas-Fort Worth Federal Statistical Research Data Center, and the support of AHRQ is acknowledged. Results and conclusions are those of the author and do not indicate concurrence by AHRQ or the Department of Health and Human Services